

# **Report on implementation of the MK-model for resonance single-pion production into GENIE**

I.D. Kakorin, K.S. Kuzmin, V.A. Naumov  
Bogoliubov Laboratory of Theoretical Physics  
Joint Institute for Nuclear Research  
Dubna

June 18, 2020


## Abstract

In this report, we intend to set out the questions and problems which have arisen during the implementation of the MK-model [1–4] in GENIE Generator.

The report consists of two parts. In the first part, we list yet unfixed bugs and mistakes, because they require thorough study for their correction. In the second part, we list resolved problems. The bugs and mistakes in this part have been fixed either after intensive discussions with the author of the model<sup>1</sup> or if the bugs are trivial.

This is a living document which is being constantly updated. Any comments, suggestions and corrections are welcome.

In the current version of this report, we discuss the latest version of the Monoo code in which she got out of the phase factors earlier belonging to the vector and axial parts of each resonance amplitude and were the subject of fine-tuning. From our point of view, this simplification (rejecting the phase factors) is a very right step.



<sup>1</sup>In sections of this part, we point the version of the erratum or code where they were fixed.

# Contents

<b>1</b>	<b>Still unfixed problems, bugs and mistakes</b>	<b>1</b>
1.1	The values of some resonance parameters are incorrect . . . . .	1
1.2	Value of $\hat{\kappa}$ -factor is incorrect . . . . .	2
1.3	Ambiguity in calculation of signs of the amplitudes . . . . .	2
1.3.1	General relations . . . . .	2
1.3.2	Derivation by direct way . . . . .	6
1.3.3	Derivation by using symmetry properties . . . . .	6
1.3.4	Paradox and probable explanation . . . . .	6
1.4	Problem with transition from neutrino to antineutrino case . . . . .	8
<b>2</b>	<b>Fixed problems, bugs and mistakes</b>	<b>18</b>
2.1	Incorrect phase factors for resonance and background amplitudes . . . . .	18
2.2	Incorrect helicity amplitudes . . . . .	18
2.3	Erroneous dynamical form factor B . . . . .	19
2.4	Mistaken cross-section formula for NC-processes in the code . . . . .	19
2.5	Incorrect $C_j$ -signs . . . . .	20
2.6	Problem with multipole expansion of some amplitudes in the code . . . . .	21
2.7	Expressions for EM-amplitudes contains mistake . . . . .	21
2.8	Error in the dynamical amplitudes for resonance $D_{13}(1520)$ . . . . .	21
2.9	Wrong sign for As-values in the code . . . . .	22
2.10	Mistype in a formula for cross section . . . . .	22
2.11	Problem with phase factors for resonance amplitudes . . . . .	22

# Chapter 1

## Still unfixed problems, bugs and mistakes

### 1.1 The values of some resonance parameters are incorrect

The values of resonance masses defined in the latest version of the file *mkcons.h* are wrong:

```
const double MR[17] = {1.232, 1.440, 1.515, 1.53, 1.57,  
                      1.61, 1.65, 1.675, 1.685, 1.72,  
                      1.71, 1.71, 1.72, 1.88, 1.9, 1.92, 1.93};
```

because they should be

```
const double MR[17] = {1.232, 1.440, 1.515, 1.530, 1.610,  
                      1.650, 1.675, 1.685, 1.720, 1.710,  
                      1.710, 1.720, 1.880, 1.900, 1.920, 1.930, 1.570};
```

according to the order in which resonance amplitudes are calculated in the files *MK\_imode1.cc*–*MK\_imode17.cc*:

```
P_{33}(1232) **** IBLOCK=0 ****  
P_{11}(1440) **** IBLOCK=1 ****  
D_{13}(1520) **** IBLOCK=2 ****  
S_{11}(1535) **** IBLOCK=3 ****  
S_{31}(1620) **** IBLOCK=4 ****  
S_{11}(1650) **** IBLOCK=5 ****  
D_{15}(1675) **** IBLOCK=6 ****  
F_{15}(1680) **** IBLOCK=7 ****  
D_{13}(1700) **** IBLOCK=8 ****  
D_{33}(1700) **** IBLOCK=9 ****  
P_{11}(1710) **** IBLOCK=10 ****  
P_{13}(1720) **** IBLOCK=11 ****  
F_{35}(1905) **** IBLOCK=12 ****  
P_{31}(1910) **** IBLOCK=13 ****  
P_{33}(1920) **** IBLOCK=14 ****  
F_{37}(1950) **** IBLOCK=15 ****  
P_{33}(1600) **** IBLOCK=16 ****
```

The same is also true for the widths and branching ratios. However such inputs as a signs of the angular Clebsch–Gordan coefficients, angular momentum and total angular momentum projections are defined correctly. Since it is not quite clear which set of resonance amplitudes is used in the latest version of the code, it becomes problematic to determine whether the values of  $\sigma^D$  is correct or not since they differ in the previous and latest code version (see also section 1.3.4).

## 1.2 Value of $\hat{\kappa}$ -factor is incorrect

Recall that the MK model is essentially based on Rein model [5]. It is obvious from comparison of Eqs. (4.15) in Ref. [1] and (40a) in Ref. [5] that the  $\kappa$ -factor should be replaced by  $\hat{\kappa}$ , which is defined after Eq. (39c) in Ref. [5].

To calculate the  $\hat{\kappa}$ -factor one needs know the ratio of isospin coefficients

$$\zeta = \frac{c_I}{a_I},$$

where  $a_I$  and  $c_I$  are defined in Eqs. (8) and (24) of Rein's paper [5]. Note, that the isospin coefficients  $a_I$  and  $c_I$ <sup>1</sup> are given in the explicit form only for several channel in Ref. [5]. It is therefore necessary to calculate them for other channels. The absence of this factor leads to very significant diversities with the original theory (we have tested this numerically).

## 1.3 Ambiguity in calculation of signs of the amplitudes

In this section, we show that one will get a contradiction in calculation of the helicity amplitudes if one try to calculate these by two different ways: directly and by using some known symmetry relations. First, we recall the general formulas.

### 1.3.1 General relations

From Eq. (3.34)<sup>2</sup> it follows that:

$$\tilde{F}_{\lambda_2, \lambda_1}^{\lambda_k(p)}(\theta, \phi) = \frac{1}{2M} \langle N\pi, \lambda_2 | e_p^\alpha J_\alpha^V | N, \lambda_1 \rangle, \quad (1.1)$$

where  $e_p$  is the gauge boson polarization,  $p = \{L, R, +, -\}$ ,  $\lambda_k(L) = -1$ ,  $\lambda_k(R) = 1$ , and  $\lambda_k(\pm) = 0$ . We will adopt the following shorter notation of the equation (1.1):

$$\tilde{F}_{\lambda_2, \lambda_1}^{\lambda_k}(\theta, \phi) = \frac{1}{2M} \langle N\pi, \lambda_2 | e_{\lambda_k}^\alpha J_\alpha^V | N, \lambda_1 \rangle, \quad (1.2)$$

---

<sup>1</sup>We want to note here, that in Rein's paper [5], the coefficient  $c_I$  for the decay channel  $\nu p \rightarrow \ell p \pi^+$  is equal to 1, while in Ref. [6] it equal to  $\sqrt{3}$ . The origin of the factor  $\sqrt{3}$  in the Rein-Sehgal paper is clear (it arises due to the isospin symmetry), but it is not clear (for us!) why it is absent in Rein's paper. Maybe somebody can clarify this question for us?

<sup>2</sup>In this section, if a certain reference is not specified, we assume that the corresponding equation has been taken from Ref. [1].

where  $\lambda_k$  denotes either the symbols  $L, R, +, -$  or a relevant number, as the context admits. Let's now express the matrix element of single pion production as product of two matrix elements: resonance production and resonance decay to nucleon-pion pair (similar to Eq. (4.18)).

$$\langle N\pi, \lambda_2 | e^\alpha J_\alpha^V | N, \lambda_1 \rangle = \langle N\pi, \lambda_2 | R, \lambda \rangle \langle R, \lambda | e^\alpha J_\alpha^V | N, \lambda_1 \rangle. \quad (1.3)$$

First, let's find the matrix element for the resonance production. It can be found from Eq. (3.17) that

$$e^\alpha J_\alpha^V = \sum_{\sigma=(-,+)} e_\sigma^\alpha J_\alpha^V = \sum_{\sigma=(-,+)} (C_{L\sigma} e_L^\alpha + C_{R\sigma} e_R^\alpha + C_\sigma e_\sigma^\alpha) J_\alpha^V. \quad (1.4)$$

Here  $\sigma = -(+)$  denotes left(right) lepton helicity.

Using Eqs. (4.5) and (4.6), the matrix element of resonance production can be written as

$$\begin{aligned} \langle R, \lambda | e^\alpha J_\alpha^V | N, \lambda_1 \rangle &= 2M_R \sum_{\sigma=(-,+)} \langle R, \lambda | (C_{L\sigma} e_L^\alpha + C_{R\sigma} e_R^\alpha + C_\sigma e_\sigma^\alpha) F_\alpha^V | N, \lambda_1 \rangle = \\ &= 2M_R \sum_{\sigma=(-,+)} \langle R, \lambda | C_{L\sigma} F_- + C_{R\sigma} F_+ + C_\sigma F_0^{(\sigma)} | N, \lambda_1 \rangle, \end{aligned} \quad (1.5)$$

where  $F_- = e_L^\alpha F_\alpha^V$ ,  $F_+ = e_R^\alpha F_\alpha^V$ ,  $F_0^{(-)} = e_-^\alpha F_\alpha^V$  and  $F_0^{(+)} = e_+^\alpha F_\alpha^V$ , at that  $F_-^\dagger = -F_+$ . The resonance production amplitudes in the RS model [6] are expressed as

$$f_{\pm|2s_z|} = \langle N, s_z \pm 1 | F_\pm | R, s_z \rangle, \quad f_{0\pm} = \langle N, s_z \pm \frac{1}{2} | F_0^{(\pm)} | R, s_z \pm \frac{1}{2} \rangle \quad (1.6)$$

We can choose the spin quantization axis in such a way that  $\lambda_1 = -s_{1z}$ ,  $\lambda_2 = -s_{2z}$ , and  $\lambda = s_{Rz}$ . Therefore,

$$\begin{aligned} \langle R, \lambda | e_L^\alpha F_\alpha^V | N_1, \lambda_1 = -(\lambda + 1) \rangle &= \langle R, \lambda | F_- | N_1, \lambda_1 = -(\lambda + 1) \rangle \\ &= \langle N_1, \lambda_1 = -(\lambda + 1) | F_-^\dagger | R, \lambda \rangle^* \\ &= \langle N_1, \lambda_1 = -(\lambda + 1) | -F_+ | R, \lambda \rangle^* \\ &= -\langle N_1, s_{1z} = (\lambda + 1) | F_+ | R, s_{Rz} = \lambda \rangle^* \\ &= -f_{+|2\lambda|}^{*V} = -f_{+|2\lambda|}^V \\ \langle R, \lambda | e_R^\alpha F_\alpha^V | N_1, \lambda_1 = -(\lambda - 1) \rangle &= \langle R, \lambda | F_+ | N_1, \lambda_1 = -(\lambda - 1) \rangle \\ &= \langle N_1, \lambda_1 = -(\lambda - 1) | F_+^\dagger | R, \lambda \rangle^* \\ &= \langle N_1, \lambda_1 = -(\lambda - 1) | -F_- | R, \lambda \rangle^* \\ &= -\langle N_1, s_{1z} = (\lambda - 1) | F_- | R, s_{Rz} = \lambda \rangle^* \\ &= -f_{-|2\lambda|}^{*V} = -f_{-|2\lambda|}^V, \end{aligned} \quad (1.7)$$

$$\begin{aligned}
\langle R, \lambda = \pm \frac{1}{2} | e_{-}^{\alpha} F_{\alpha}^V | N_1, \lambda_1 = \mp \frac{1}{2} \rangle &= \langle R, \lambda = \pm \frac{1}{2} | F_0^{(-)} | N_1, \lambda_1 = \mp \frac{1}{2} \rangle \\
&= \langle N_1, \lambda_1 = \mp \frac{1}{2} | F_0^{(-)\dagger} | R, \lambda = \pm \frac{1}{2} \rangle^* \\
&= \langle N_1, \lambda_1 = \mp \frac{1}{2} | F_0^{(-)} | R, \lambda = \pm \frac{1}{2} \rangle^* \\
&= \langle N_1, s_{1z} = \pm \frac{1}{2} | F_0^{(-)} | R, s_{Rz} = \pm \frac{1}{2} \rangle^* \\
&= f_{0\pm}^{*V(-)} = f_{0\pm}^{V(-)}, \\
\langle R, \lambda = \pm \frac{1}{2} | e_{+}^{\alpha} F_{\alpha}^V | N_1, \lambda_1 = \mp \frac{1}{2} \rangle &= \langle R, \lambda = \pm \frac{1}{2} | F_0^{(+)} | N_1, \lambda_1 = \mp \frac{1}{2} \rangle \\
&= \langle N_1, \lambda_1 = \mp \frac{1}{2} | F_0^{(+)\dagger} | R, \lambda = \pm \frac{1}{2} \rangle^* \\
&= \langle N_1, \lambda_1 = \mp \frac{1}{2} | F_0^{(+)} | R, \lambda = \pm \frac{1}{2} \rangle^* \\
&= \langle N_1, s_{1z} = \pm \frac{1}{2} | F_0^{(+)} | R, s_{Rz} = \pm \frac{1}{2} \rangle^* \\
&= f_{0\pm}^{*V(+)} = f_{0\pm}^{V(+)}.
\end{aligned} \tag{1.8}$$

The resonance decay matrix element is given in Eq. (4.20):

$$\langle N\pi, \lambda_2 | R, \lambda \rangle = \sigma^D C_{N\pi}^j \sqrt{\chi_E \kappa} C_{N\pi}^I f_{BW} \tag{1.9}$$

and the meaning of all symbols can be found in Ref. [1].

The relation between the standard helicity amplitudes and helicity amplitudes defined by Eq. (1.2) is given by Eq. (3.67):

$$F_{\mu, \lambda}(\theta, \phi) = e^{i[\lambda_1 \pi + \lambda_2(\pi + 2\phi)]} \tilde{F}_{\lambda_2, \lambda_1}^{\lambda_k}(\theta, \phi), \tag{1.10}$$

where  $\lambda$  is spin of resonance,  $\lambda = \lambda_k - \lambda_1$  ( $\lambda \in \{-\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{3}{2}\}$ ) and  $\mu = -\lambda_2$  ( $\mu \in \{-\frac{1}{2}, \frac{1}{2}\}$ ).

The multipole expansion of  $F_{\mu, \lambda}(\theta, \phi)$ :

$$F_{\mu, \lambda}(\theta, \phi) = \sum_j \left\{ \begin{array}{l} F_{\mu, \lambda}^j, \quad \text{if } \lambda_k \text{ is L or R} \\ F_{\mu, \lambda}^{0j}, \quad \text{if } \lambda_k \text{ is + or -} \end{array} \right\} (2j+1) d_{\lambda, \mu}^j(\theta) e^{i(\lambda - \mu)\phi}. \tag{1.11}$$

By comparing Eqs. (1.10) and (1.11) with Eqs. (4.15) and (4.26) one obtains:

$$\tilde{F}_{\lambda_2, \lambda_1}^{\lambda_k}(\theta, \phi) = (-1)^{(\lambda_1 + \lambda_2)} \sum_j \left\{ \begin{array}{l} F_{\mu, \lambda}^j, \quad \text{if } \lambda_k \text{ is L or R} \\ F_{\mu, \lambda}^{0j}, \quad \text{if } \lambda_k \text{ is + or -} \end{array} \right\} \frac{2j+1}{\sqrt{2}} d_{\lambda, \mu}^j(\theta) e^{i(\lambda_k - \lambda_1 - \lambda_2)\phi}, \tag{1.12}$$

where the values  $F_{\mu, \lambda}^j$  ( $F_{\mu, \lambda}^{0j}$ ) are the coefficients of multipole expansion. The amplitudes  $F_{\mu, \lambda}^j$  are defined similarly to Eq. (30) in Ref. [5] (except when the common phase is chosen to be +1). The superscript "0" in Eqs. (1.11) and (1.12) hereinafter is omitted and implicitly implied where it is needed. The standard helicity amplitudes possess the following symmetry properties:

$$F_{-\mu, -\lambda}(\theta, \phi) = -e^{i(\lambda - \mu)(\pi - 2\phi)} F_{\mu, \lambda}(\theta, \phi). \tag{1.13}$$

Substituting Eq. (1.11) into Eq. (1.13) we obtain

$$\begin{aligned} F_{-\mu,-\lambda}(\theta, \phi) &= -e^{i(\lambda-\mu)(\pi-2\phi)} \sum_j F_{\mu,\lambda}^j (2j+1) d_{\lambda,\mu}^j(\theta) e^{i(\lambda-\mu)\phi} \\ &= -e^{i(\lambda-\mu)\pi} \sum_j F_{\mu,\lambda}^j (2j+1) d_{\lambda,\mu}^j(\theta) e^{-i(\lambda-\mu)\phi}. \end{aligned} \quad (1.14)$$

Using the symmetry properties of the functions  $d_{\lambda\mu}^j$  (see Eq. (A1) in Ref. [7])

$$d_{\lambda,\mu}^j(\theta) = d_{-\mu,-\lambda}^j(\theta) = (-1)^{\lambda-\mu} d_{\mu,\lambda}^j(\theta), \quad (1.15)$$

one obtains

$$d_{\lambda,\mu}^j(\theta) = d_{-\mu,-\lambda}^j(\theta) = (-1)^{\lambda-\mu} d_{-\lambda,-\mu}^j(\theta). \quad (1.16)$$

Then Eq. (1.14) can be rewritten as

$$F_{-\mu,-\lambda}(\theta, \phi) = - \sum_j F_{\mu,\lambda}^j (2j+1) d_{-\lambda,-\mu}^j(\theta) e^{-i(\lambda-\mu)\phi}. \quad (1.17)$$

From the other hand, if one put  $-\mu$  and  $-\lambda$  instead of  $\mu$  and  $\lambda$  in Eq. (1.11)

$$F_{-\mu,-\lambda}(\theta, \phi) = \sum_j F_{-\mu,-\lambda}^j (2j+1) d_{-\lambda,-\mu}^j(\theta) e^{-i(\lambda-\mu)\phi}. \quad (1.18)$$

Subtracting Eq. (1.18) from Eq. (1.17) we find:

$$0 = \sum_j (F_{-\mu,-\lambda}^j + F_{\mu,\lambda}^j) (2j+1) d_{-\lambda,-\mu}^j(\theta) e^{-i(\lambda-\mu)\phi}. \quad (1.19)$$

Now let's use the orthonormality of the functions

$$\int_0^\pi d_{\lambda,\mu}^j(\theta) d_{\lambda,\mu}^{j'}(\theta) d \cos(\theta) = \frac{2}{2j+1} \delta_{jj'}. \quad (1.20)$$

Multiplying Eq. (1.18) by  $d_{\lambda,\mu}^{j'}$  and integrating, we obtain

$$F_{-\mu,-\lambda}^j = -F_{\mu,\lambda}^j. \quad (1.21)$$

The following expressions are valid:

$$\begin{aligned} \text{for } j = l + \frac{1}{2}: \quad f_{+1,+3,0+}^V &= -f_{-1,-3,0-}^V, \quad f_{+1,+3,0+}^A = f_{-1,-3,0-}^A, \\ \text{for } j = l - \frac{1}{2}: \quad f_{+1,+3,0+}^V &= f_{-1,-3,0-}^V, \quad f_{+1,+3,0+}^A = -f_{-1,-3,0-}^A. \end{aligned} \quad (1.22)$$

We have prepared everything needed to make calculation by both mentioned ways. Let's consider, for example, the calculation of  $\widetilde{F}_{\frac{1}{2}\frac{1}{2}}^L(\theta, \phi)$  for which  $\lambda_2 = \frac{1}{2}$ ,  $\lambda_1 = \frac{1}{2}$ ,  $\lambda_k = -1$ ,  $\mu = -\frac{1}{2}$  and  $\lambda = -\frac{3}{2}$ .



### 1.3.2 Derivation by direct way

Using Eqs. (1.2)–(1.9) and (1.12) we can find the coefficients of multipole expansion for  $\tilde{F}_{\frac{1}{2}, \frac{1}{2}}^{-1}(\theta, \phi)$ :

$$\begin{aligned}
F_{-\frac{1}{2}, -\frac{3}{2}}^j &= \frac{1}{2M} \langle N\pi, \lambda_2 = \frac{1}{2} | e_L^\alpha J_\alpha^V | N, \lambda_1 = \frac{1}{2} \rangle \\
&= \frac{1}{2M} \langle N\pi, \lambda_2 = \frac{1}{2} | R, \lambda = -\frac{3}{2} \rangle \langle R, \lambda = -\frac{3}{2} | e_L^\alpha J_\alpha^V | N, \lambda_1 = \frac{1}{2} \rangle \\
&= \frac{M_R}{M} \sigma^D C_{N\pi}^j \sqrt{\chi_E} \kappa C_{N\pi}^I f_{BW}(-f_{+3}^V).
\end{aligned} \tag{1.23}$$

### 1.3.3 Derivation by using symmetry properties

Using the symmetry property (1.21) for the multipole expansion coefficients, one obtains:

$$F_{-\frac{1}{2}, -\frac{3}{2}}^j = -F_{\frac{1}{2}, \frac{3}{2}}^j. \tag{1.24}$$

The values  $F_{\frac{1}{2}, \frac{3}{2}}^j$  are multipole expansion coefficients for the amplitude  $\tilde{F}_{-\frac{1}{2}, -\frac{1}{2}}^{+1}(\theta, \phi)$ . Then

$$\begin{aligned}
F_{-\frac{1}{2}, -\frac{3}{2}}^j &= -\frac{1}{2M} \langle N\pi, \lambda_2 = -\frac{1}{2} | R, \lambda = \frac{3}{2} \rangle \langle R, \lambda = \frac{3}{2} | e_R^\alpha J_\alpha^V | N, \lambda_1 = -\frac{1}{2} \rangle \\
&= \frac{M_R}{M} \sigma^D C_{N\pi}^j \sqrt{\chi_E} \kappa C_{N\pi}^I f_{BW} f_{-3}^V.
\end{aligned} \tag{1.25}$$

### 1.3.4 Paradox and probable explanation

Equations (1.23) and (1.25) are consistent only for the resonances with  $j = l + \frac{1}{2}$  (for which  $f_{-3}^V = -f_{+3}^V$ ) as it can be seen from Eq. (1.22). But there is evident **contradiction** for the resonances with  $j = l - \frac{1}{2}$  (for which  $f_{-3}^V = +f_{+3}^V$ ).

What is the reason of such contradiction? We understand that in the case of  $j = l - \frac{1}{2}$  one must take into account the signs of  $C_{N\pi}^j$  and  $\sigma^D$ , but the above formulas were derived for arbitrary  $C_{N\pi}^j$  and  $\sigma^D$ ; so there is some other nuance which we cannot find. This is very important because from our calculation (see bellow) we obtain the set of signs which is in general **different** from ones in Refs. [1, 3]. And this muddle almost definitely sits in the above paradox.

We think that the problem is with the symmetry condition stated by Eqs. (15a) and (15b) in Ref. [5], namely, they are only valid for the resonances with  $j = l + \frac{1}{2}$ , but for the case  $j = l - \frac{1}{2}$  one needs to use different relations with opposite signs. Some hints on this can be seen from Eq. (44) in Ref. [7] and in the words after Eq. (18f) in Ref. [5], but this question requires a more thorough study.

We calculated all amplitudes by the direct method, because it does not lead, in our opinion, to any contradictions, and we present the result in Table 1.1. The latest versions of the amplitudes calculated by Minoo are presented in Table III of erratum for Ref. [3]. One can see that the amplitudes differ from those calculated by the direct method.

It is difficult to understand which set of the amplitude signs is used in Minoo's code. The problem is that the signs belonging to one terms is sometimes assigned to another term. Considering numerous versions that increased as a result of several corrections, the problem with

signs becomes even more complicated than it was before corrections. Several examples of the problem with signs:

- the sign of the amplitude itself, which depends on whether  $j$  is equal to  $l + \frac{1}{2}$  or  $l - \frac{1}{2}$ ;
- the sign of the functions  $d_{\lambda,\mu}^j$ , because Minoo coded only  $d_{\lambda,\mu}^j$  with  $\lambda > 0$  and then she finds other values using the symmetry properties of the  $d$ -functions (1.15) and (1.16);
- the sign of  $\sigma^D$ , which sits in the term  $\mathcal{D}^j(R)$  (see Eq.(25) of Ref. [3]) and is different from the original Rein's signs <sup>3</sup> (see Table 5.3 in Ref. [1]).

However, one thing is obvious: the amplitude signs in the code definitely differ from those in the erratum [3]. Let's note that it is impossible to control (adjust) the signs by comparison of calculations with experimental data since currently available data are very slowly sensible to these signs. But interference between the high-mass resonances is very responsible to them. This will be important in the future experiments.

Therefore we use the signs as is in Minoo's code for now. However it poses a problem as it is impossible to check whether the signs are correct (see also sections 1.1 and 2.6).

$\lambda_2$	$\lambda_1$	$\tilde{F}_{\lambda_2,\lambda_1}^L(\theta, \phi) - \tilde{G}_{\lambda_2,\lambda_1}^L(\theta, \phi)$	$\tilde{F}_{\lambda_2,\lambda_1}^R(\theta, \phi) - \tilde{G}_{\lambda_2,\lambda_1}^R(\theta, \phi)$
$\frac{1}{2}$	$\frac{1}{2}$	$-\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{+3}(R) d_{\frac{3}{2}\frac{1}{2}}^j(\theta) e^{-2i\phi}$	$+\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{-1}(R) d_{\frac{1}{2}-\frac{1}{2}}^j(\theta)$
$-\frac{1}{2}$	$\frac{1}{2}$	$+\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{+3}(R) d_{\frac{3}{2}-\frac{1}{2}}^j(\theta) e^{-i\phi}$	$+\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{-1}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{i\phi}$
$\frac{1}{2}$	$-\frac{1}{2}$	$+\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{+1}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{-i\phi}$	$+\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{-3}(R) d_{\frac{3}{2}-\frac{1}{2}}^j(\theta) e^{i\phi}$
$-\frac{1}{2}$	$-\frac{1}{2}$	$-\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{+1}(R) d_{\frac{1}{2}-\frac{1}{2}}^j(\theta)$	$+\sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{-3}(R) d_{\frac{3}{2}\frac{1}{2}}^j(\theta) e^{2i\phi}$
		$\tilde{F}_{\lambda_2,\lambda_1}^-(\theta, \phi) - \tilde{G}_{\lambda_2,\lambda_1}^-(\theta, \phi)$	$\tilde{F}_{\lambda_2,\lambda_1}^+(\theta, \phi) - \tilde{F}_{\lambda_2,\lambda_1}^+(\theta, \phi)$
$\frac{1}{2}$	$\frac{1}{2}$	$+\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0-}^{(-)}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{-i\phi}$	$+\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0-}^{(+)}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{-i\phi}$
$-\frac{1}{2}$	$\frac{1}{2}$	$-\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0-}^{(-)}(R) d_{\frac{1}{2}-\frac{1}{2}}^j(\theta)$	$-\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0-}^{(+)}(R) d_{\frac{1}{2}-\frac{1}{2}}^j(\theta)$
$\frac{1}{2}$	$-\frac{1}{2}$	$+\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0+}^{(-)}(R) d_{\frac{1}{2}-\frac{1}{2}}^j(\theta)$	$+\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0+}^{(+)}(R) d_{\frac{1}{2}-\frac{1}{2}}^j(\theta)$
$-\frac{1}{2}$	$-\frac{1}{2}$	$+\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0+}^{(-)}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{i\phi}$	$+\frac{ \mathbf{k} }{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0+}^{(+)}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{i\phi}$

Table 1.1: Vector helicity amplitudes of resonant interactions calculated by direct method.

<sup>3</sup>The sign of resonance  $P_{11}(1710)$ ,  $\sigma^D$ , given in Ref. [6] is equal to "+". It was changed in Ref. [5] and is now equal to "-".

## 1.4 Problem with transition from neutrino to antineutrino case

Since this problem is one of the key points, we consider it in some detail. Let's start with a reminder of several basic formulas derived in Ref. [8]. The components of the leptonic current with the lepton helicity  $\lambda$  measured in lab. frame, can be written in the resonance rest frame (RRF)<sup>4</sup> as

$$\begin{aligned}
 j_0^* &= N_\lambda m_\ell \frac{E_\nu}{W} (1 - \lambda \cos \theta) (M - E_\ell - \lambda P_\ell), \\
 j_x^* &= N_\lambda m_\ell \frac{E_\nu}{|\mathbf{q}|} \sin \theta (P_\ell - \lambda E_\nu), \\
 j_y^* &= i \lambda N_\lambda m_\ell E_\nu \sin \theta, \\
 j_z^* &= N_\lambda m_\ell \frac{E_\nu}{|\mathbf{q}| W} (1 - \lambda \cos \theta) [(E_\nu + \lambda P_\ell) (M - E_\ell) + P_\ell (\lambda E_\nu + 2E_\nu \cos \theta - P_\ell)].
 \end{aligned} \tag{1.26}$$

Here  $E_\nu$ ,  $E_\ell$ ,  $\theta$ , and  $m_\ell$  are, respectively, the incident neutrino energy, lepton energy, scattering angle, and mass of the lepton,  $P_\ell = \sqrt{E_\ell^2 - m_\ell^2}$ ,  $\mathbf{q} = \mathbf{k}_\nu - \mathbf{k}_\ell$ , the vectors  $\mathbf{k}_\nu$  and  $\mathbf{k}_\ell$  are the 3-momenta of the neutrino and lepton. All variables are written in lab. frame. The normalization constant  $N_\lambda$  is expressed in terms of the kinematic variables,

$$N_\lambda = \frac{1}{m_\ell} \sqrt{\frac{E_\ell \mp \lambda P_\ell}{E_\nu (1 \mp \lambda \cos \theta)}}, \tag{1.27}$$

where the upper (lower) sign is for  $\nu_\ell$  ( $\bar{\nu}_\ell$ ).

On the other hand, in the spirit of the RS model, the leptonic current can be treated as the intermediate  $W$  boson polarization 4-vector and may be decomposed into three polarization 4-vectors  $e_L$ ,  $e_R$ , and  $e_S(\lambda) \equiv e_{(\lambda)}$  corresponding to left-handed, right-handed and scalar polarizations:

$$j_\lambda^\alpha = K^{-1} [c_L^\lambda e_L^\alpha + c_R^\lambda e_R^\alpha + c_S^\lambda e_{(\lambda)}^\alpha], \tag{1.28}$$

$$e_L^\alpha = \frac{1}{\sqrt{2}} (0, 1, -i, 0),$$

$$e_R^\alpha = \frac{1}{\sqrt{2}} (0, -1, -i, 0), \tag{1.29}$$

$$e_{(\lambda)}^\alpha = \frac{1}{\sqrt{Q^2}} (Q_{(\lambda)}^*, 0, 0, \nu_{(\lambda)}^*).$$

Note that the 4-vectors  $e_L$  and  $e_R$  are exactly the same as in the RS model [6], while  $e_{(\lambda)}$  has been modified to include the lepton mass effect; its components are given by

$$Q_{(\lambda)}^* = \frac{K \sqrt{Q^2}}{c_S^\lambda} j_0^*, \quad \nu_{(\lambda)}^* = \frac{K \sqrt{Q^2}}{c_S^\lambda} j_z^*, \quad K = \frac{|\mathbf{q}|}{E_\nu \sqrt{2Q^2}}. \tag{1.30}$$

The polarization vectors form an orthonormal set:

$$e_i^\alpha e_{j\alpha} = \delta_{ij}, \quad i, j = L, R, S.$$

<sup>4</sup>Here and below we mark this frame with asterisk (\*).

Taking this into account, the coefficients  $c_i^\lambda$  are explicitly defined through the components  $j_\alpha^\lambda$  in RRF as

$$c_L^\lambda = \frac{K}{\sqrt{2}} (j_x^* + ij_y^*), \quad c_R^\lambda = -\frac{K}{\sqrt{2}} (j_x^* - ij_y^*), \quad c_S^\lambda = K \sqrt{|(j_0^*)^2 - (j_z^*)^2|}. \quad (1.31)$$

The sign in the last relation is chosen to be positive, by definition. The currents for neutrino and antineutrino can be related with Eqs. (2) and (3) in Ref. [6]

$$\bar{j}_\alpha^\lambda = -\lambda (j_\alpha^{-\lambda})^*. \quad (1.32)$$

This identity is an intrinsic property of the leptonic current and does not depend on the nature of the process in which it participates. By using Eqs. (1.26) and (1.32) we can write

$$\begin{aligned} [j_0^{\star\lambda}]_{\bar{\nu}} &= -\lambda [j_0^{\star-\lambda}]_{\nu}, \\ [j_x^{\star\lambda}]_{\bar{\nu}} &= -\lambda [j_x^{\star-\lambda}]_{\nu}, \\ [j_y^{\star\lambda}]_{\bar{\nu}} &= +\lambda [j_y^{\star+\lambda}]_{\nu}, \\ [j_z^{\star\lambda}]_{\bar{\nu}} &= -\lambda [j_z^{\star+\lambda}]_{\nu}. \end{aligned} \quad (1.33)$$

The identities (1.31) and (1.33) allow us to derive relations between the coefficients  $c_L$ ,  $c_R$  and  $c_S$  for the neutrino and antineutrino cases:

$$\begin{aligned} [c_L^\lambda]_{\bar{\nu}} &= \frac{K}{\sqrt{2}} \left( [j_x^{\star\lambda}]_{\bar{\nu}} + i [j_y^{\star\lambda}]_{\bar{\nu}} \right) = \frac{K}{\sqrt{2}} \left( -\lambda [j_x^{\star-\lambda}]_{\nu} + i\lambda [j_y^{\star+\lambda}]_{\nu} \right) = \lambda [c_R^{-\lambda}]_{\nu}, \\ [c_R^\lambda]_{\bar{\nu}} &= -\frac{K}{\sqrt{2}} \left( [j_x^{\star\lambda}]_{\bar{\nu}} - i [j_y^{\star\lambda}]_{\bar{\nu}} \right) = -\frac{K}{\sqrt{2}} \left( -\lambda [j_x^{\star-\lambda}]_{\nu} - i\lambda [j_y^{\star+\lambda}]_{\nu} \right) = \lambda [c_L^{-\lambda}]_{\nu}, \\ [c_S^\lambda]_{\bar{\nu}} &= K \sqrt{|(j_0^{\star\lambda})_{\bar{\nu}}^2 - (j_z^{\star\lambda})_{\bar{\nu}}^2|} = K \sqrt{|(-\lambda [j_0^{\star-\lambda}]_{\nu})^2 - (-\lambda [j_z^{\star+\lambda}]_{\nu})^2|} \\ &= \pm \lambda K \sqrt{|(j_0^{\star-\lambda})_{\nu}^2 - (j_z^{\star+\lambda})_{\nu}^2|} = \pm \lambda [c_S^{-\lambda}]_{\nu}. \end{aligned}$$

Considering that

$$\begin{aligned} K^{-1} [c_S^\lambda]_{\bar{\nu}} [e_{(\lambda)}^\alpha]_{\bar{\nu}} &= ([j_0^{\star\lambda}]_{\bar{\nu}}, 0, 0, [j_z^{\star\lambda}]_{\bar{\nu}}) = -\lambda ([j_0^{\star-\lambda}]_{\nu}, 0, 0, [j_z^{\star+\lambda}]_{\nu}) \\ &= -\lambda K^{-1} [c_S^{-\lambda}]_{\nu} [e_{(-\lambda)}^\alpha]_{\nu}, \end{aligned}$$

we obtain

$$[c_S^\lambda]_{\bar{\nu}} [e_{(\lambda)}^\alpha]_{\bar{\nu}} = -\lambda [c_S^{-\lambda}]_{\nu} [e_{(-\lambda)}^\alpha]_{\nu}. \quad (1.34)$$

To derive the exact relation for the coefficient  $c_S^\lambda$ , let's consider the limiting case of a massless final-state lepton. Since the massless lepton helicity can only be -1 (for incoming neutrino) or +1 (for incoming antineutrino), we have

$$\left. \begin{aligned} [\mathcal{Q}_{(-1)}^*]_{\nu} &\rightarrow |\mathbf{q}^*| \quad \text{and} \quad [\nu_{(-1)}^*]_{\nu} \rightarrow \nu^* \\ [\mathcal{Q}_{(+1)}^*]_{\bar{\nu}} &\rightarrow |\mathbf{q}^*| \quad \text{and} \quad [\nu_{(+1)}^*]_{\bar{\nu}} \rightarrow \nu^* \end{aligned} \right\} \text{ as } m_\ell \rightarrow 0.$$

This implies

$$[\mathcal{Q}_{(-1)}^*]_{\nu} \rightarrow [\mathcal{Q}_{(+1)}^*]_{\bar{\nu}} \quad \text{and} \quad [\nu_{(-1)}^*]_{\nu} \rightarrow [\nu_{(+1)}^*]_{\bar{\nu}} \quad \text{as } m_\ell \rightarrow 0. \quad (1.35)$$

If the equality  $[c_S^\lambda]_{\bar{\nu}} = \lambda [c_S^{-\lambda}]_\nu$  is valid then, according to Eq. (1.34),

$$[e_{(\lambda)}^\alpha]_{\bar{\nu}} = -[e_{(-\lambda)}^\alpha]_\nu. \quad (1.36)$$

In the opposite case, from the equality  $[c_S^\lambda]_{\bar{\nu}} = -\lambda [c_S^{-\lambda}]_\nu$  it follows that

$$[e_{(\lambda)}^\alpha]_{\bar{\nu}} = [e_{(-\lambda)}^\alpha]_\nu. \quad (1.37)$$

Since Eq. (1.36) contradicts the conditions (1.35), only the equality  $[c_S^\lambda]_{\bar{\nu}} = -\lambda [c_S^{-\lambda}]_\nu$  and Eq. (1.37) is correct, from which it follows that

$$[Q_{(\lambda)}^*]_{\bar{\nu}} = [Q_{(-\lambda)}^*]_\nu, \quad [\nu_{(\lambda)}^*]_{\bar{\nu}} = [\nu_{(-\lambda)}^*]_\nu.$$

As a result, we get

$$[Q_{(\lambda)}^*]_{\bar{\nu}} = [Q_{(-\lambda)}^*]_\nu, \quad [\nu_{(\lambda)}^*]_{\bar{\nu}} = [\nu_{(-\lambda)}^*]_\nu, \quad (1.38)$$

$$[c_L^\lambda]_{\bar{\nu}} = \lambda [c_R^{-\lambda}]_\nu, \quad [c_R^\lambda]_{\bar{\nu}} = \lambda [c_L^{-\lambda}]_\nu, \quad [c_S^\lambda]_{\bar{\nu}} = -\lambda [c_S^{-\lambda}]_\nu. \quad (1.39)$$

We want to stress that Eqs. (1.38) and (1.39) were obtained by using only the properties of the leptonic current and do not depend on features of the hadronic current.

Minoo works in the similar framework. The differences between her notations and those used in Ref. [8] are:

- leptonic current denoted as  $\varepsilon_\lambda^\alpha$  is equal to  $2j_\alpha^\lambda$  (compare Eq. (1) in Ref. [8] and Eq. (5) in Ref. [3]) and is expressed in terms of kinematic variables defined in RRF, while in Ref. [8] it is expressed through the lab. frame kinematic variables;
- The coefficients  $c_L^\lambda$ ,  $c_R^\lambda$ , and  $c_S^\lambda$  used in Ref. [8] are denoted in Minoo's paper as  $C_{L\lambda}$ ,  $C_{R\lambda}$ , and  $C_\lambda$ , respectively;
- the polarization vectors  $e_L^\alpha$ ,  $e_R^\alpha$ , and  $e_\lambda^\alpha$  used in Ref. [8] are denoted as, respectively,  $e_L^\alpha$ ,  $e_R^\alpha$ , and  $e_\lambda^\alpha$  in Minoo's paper.

The corresponding values defined in Minoo's paper [3] are following:

$$\begin{aligned} e_L^\alpha &= \frac{1}{\sqrt{2}} (0, 1, -i, 0), \\ e_R^\alpha &= \frac{1}{\sqrt{2}} (0, -1, -i, 0), \\ e_\lambda^\alpha &= \frac{1}{\sqrt{|(\varepsilon_\lambda^0)^2 - (\varepsilon_\lambda^3)^2|}} (\varepsilon_\lambda^0, 0, 0, \varepsilon_\lambda^3). \end{aligned} \quad (1.40)$$

and

$$\begin{aligned} C_{L\lambda} &= \frac{1}{\sqrt{2}} (\varepsilon_\lambda^1 + i\varepsilon_\lambda^2), \\ C_{R\lambda} &= -\frac{1}{\sqrt{2}} (\varepsilon_\lambda^1 - i\varepsilon_\lambda^2), \\ C_\lambda &= \sqrt{|(\varepsilon_\lambda^0)^2 - (\varepsilon_\lambda^3)^2|}. \end{aligned} \quad (1.41)$$

It can be seen that the values  $e_L^\alpha$  and  $e_R^\alpha$  from Refs. [3] and [8] are identical. Taking into account Eqs. (1.28)–(1.31), (1.40), (1.41) and  $\varepsilon_\lambda^\alpha = 2j_\lambda^\alpha$  we can relate the corresponding values:

$$\begin{aligned}
C_{L_\lambda} e_L^\alpha + C_{R_\lambda} e_R^\alpha + C_\lambda e_\lambda^\alpha &= 2K^{-1} [c_L^\lambda e_L^\alpha + c_R^\lambda e_R^\alpha + c_S^\lambda e_{(\lambda)}^\alpha] \\
\Downarrow \\
C_{L_\lambda} &= 2K^{-1} c_L^\lambda, \quad C_{R_\lambda} = 2K^{-1} c_R^\lambda, \quad C_\lambda e_\lambda^\alpha = 2K^{-1} c_S^\lambda e_{(\lambda)}^\alpha, \\
C_\lambda &= \sqrt{|(\varepsilon_\lambda^0)^2 - (\varepsilon_\lambda^3)^2|} = 2K^{-1} K \sqrt{|(j_0^*)^2 - (j_z^*)^2|} = 2K^{-1} c_S^\lambda \\
\Downarrow \\
e_\lambda^\alpha &= e_{(\lambda)}^\alpha.
\end{aligned}$$

Thus we obtained

$$C_{L_\lambda} = 2K^{-1} c_L^\lambda, \quad C_{R_\lambda} = 2K^{-1} c_R^\lambda, \quad C_\lambda = 2K^{-1} c_S^\lambda, \quad e_\lambda^\alpha = e_{(\lambda)}^\alpha. \quad (1.42)$$

Therefore the relation similar to (1.39) must be hold:

$$[C_{L_\lambda}]_{\bar{\nu}} = \lambda [C_{R_{-\lambda}}]_{\nu}, \quad [C_{R_\lambda}]_{\bar{\nu}} = \lambda [C_{L_{-\lambda}}]_{\nu}, \quad [C_{S_\lambda}]_{\bar{\nu}} = -\lambda [C_{S_{-\lambda}}]_{\nu}. \quad (1.43)$$

For example, let us prove the first relation:

$$[C_{L_\lambda}]_{\bar{\nu}} = 2K^{-1} [c_L^\lambda]_{\bar{\nu}} = 2K^{-1} \lambda [c_R^{-\lambda}]_{\nu} = \lambda [C_{R_{-\lambda}}]_{\nu}.$$

However, in Minoo's paper it is proposed to use different relation (see words after Eq. (18) in Ref. [3] or words after Eq. (3.60) in Ref. [2]):

$$[C_{L_\lambda}]_{\bar{\nu}} = [C_{R_\lambda}]_{\nu}, \quad [C_{R_\lambda}]_{\bar{\nu}} = [C_{L_\lambda}]_{\nu}, \quad [C_{S_\lambda}]_{\bar{\nu}} = [C_{S_\lambda}]_{\nu}. \quad (1.44)$$

The last conditions can be rewritten in terms of Ref. [8] as follows

$$[c_L^\lambda]_{\bar{\nu}} = [c_R^\lambda]_{\nu}, \quad [c_R^\lambda]_{\bar{\nu}} = [c_L^\lambda]_{\nu}, \quad [c_S^\lambda]_{\bar{\nu}} = [c_S^\lambda]_{\nu}, \quad (1.45)$$

and one should apply this "recipe" to final formula for the differential cross section. This means that one can do all calculation exactly as as for the neutrino case and then replace the coefficients  $c_i^\lambda$  for neutrino by ones for antineutrino.

Let's see what this leads to. First, we study whether there is a difference between the double-differential cross section  $d\sigma/dWdQ^2$  and the polarization density matrix

$$\rho = \frac{1}{2} (1 + \sigma \mathcal{P})$$

obtained by using the relations (1.39) and (1.45) (or (1.43) and (1.44)) in the formalism developed in Ref. [8]. Again let us remind all needed points here.

The elements of the polarization matrix are given by following formulas (see Ref. [8]):

$$\rho_{\lambda\lambda'} = \frac{\Sigma_{\lambda\lambda'}}{\Sigma_{++} + \Sigma_{--}}, \quad \Sigma_{\lambda\lambda'} = \sum_{i=L,R,S} c_i^\lambda c_i^{\lambda'} \sigma_i^{\lambda\lambda'}, \quad (1.46)$$

and the differential cross section of unpolarized lepton production is given by

$$\frac{d^2\sigma}{dQ^2 dW^2} = \frac{G_F^2 \cos^2 \theta_C Q^2}{2\pi^2 M |\mathbf{q}|^2} (\Sigma_{++} + \Sigma_{--}). \quad (1.47)$$

The partial cross sections are found to be the bilinear superpositions of the reduced amplitudes for producing a  $N\pi$  final state with allowed isospin by a charged isovector current:

$$\sigma_{L,R}^{\lambda\lambda'} = \frac{\pi W}{2M} \left( A_{\pm 3}^{\lambda} A_{\pm 3}^{\lambda'} + A_{\pm 1}^{\lambda} A_{\pm 1}^{\lambda'} \right), \quad (1.48)$$

$$\sigma_S^{\lambda\lambda'} = \frac{\pi M |\mathbf{q}|^2}{2WQ^2} \left( A_{0+}^{\lambda} A_{0+}^{\lambda'} + A_{0-}^{\lambda} A_{0-}^{\lambda'} \right). \quad (1.49)$$

The amplitudes for neutrino induced reactions are

$$A_{\varkappa}^{\lambda} (p\pi^+) = \sqrt{3} \sum_{(I=3/2)} a_{\varkappa}^{\lambda} (N_3^*), \quad (1.50)$$

$$A_{\varkappa}^{\lambda} (p\pi^0) = \sqrt{\frac{2}{3}} \sum_{(I=3/2)} a_{\varkappa}^{\lambda} (N_3^*) - \sqrt{\frac{1}{3}} \sum_{(I=1/2)} a_{\varkappa}^{\lambda} (N_1^*), \quad (1.51)$$

$$A_{\varkappa}^{\lambda} (n\pi^+) = \sqrt{\frac{1}{3}} \sum_{(I=3/2)} a_{\varkappa}^{\lambda} (N_3^*) + \sqrt{\frac{2}{3}} \sum_{(I=1/2)} a_{\varkappa}^{\lambda} (N_1^*). \quad (1.52)$$

Here  $\varkappa = \pm 3, \pm 1, 0\pm$  and only those resonances are allowed to interfere which have the same spin and orbital angular momentum.

Any amplitude  $a_{\varkappa}^{\lambda} (N_i^*)$  referring to one single resonance  $N_i^*$  in a definite state of isospin, charge and helicity consists of two factors which describe the production and subsequent decay of the resonance:

$$a_{\varkappa}^{\lambda} (N_i^*) = f_{\varkappa}^{\lambda} (\nu N \rightarrow N_i^*) \eta (N_i^* \rightarrow N\pi) \equiv f_{\varkappa}^{\lambda(i)} \eta^{(i)}.$$

The decay amplitudes,  $\eta^{(i)}$ , can be split into three factors,

$$\eta^{(i)} = \text{sign}(N_i^*) \sqrt{\chi_i} \eta_{\text{BW}}^{(i)}(W),$$

irrespective of isospin, charge or helicity of the resonance. Here, the first factor is the decay sign for resonance  $N_i^*$  (see Table III of Ref. [6]),  $\chi_i$  is the elasticity of the resonance taking care of the branching ratio into the  $\pi N$  final state and  $\eta_{\text{BW}}^{(i)}(W)$  is the properly normalized Breit-Wigner term with the running width specified by the  $\pi N$  partial wave from which the resonance arises (Eq. (2.31) in Ref. [6]).

The resonance production amplitudes,  $f_{\varkappa}^{\lambda(i)}$ , can be calculated within the FKR quark model in exactly the same way as in Ref. [6]. It can be shown that they have the same structure as that given in Table II of Ref. [6] with the only important difference: the three coefficient functions  $S$ ,  $B$  and  $C$  involved into the definitions of the amplitudes have to be modified. Indeed, since the structure of the polarization 4-vector  $e_{(\lambda)}^{\alpha}$  has been changed with respect to that of the original RS model (by including the lepton mass and spin), we have to recalculate its inner products with the vector and axial hadronic currents. To do this, we used the explicit form for the FKR currents given by Ravndal [9]. As a result, the coefficients  $S$ ,  $B$  and  $C$  (and thus the resonance production amplitudes) become parametrically dependent of the lepton mass and helicity:

$$S_{(\lambda)} = S_{(\lambda)}^V = (\nu_{(\lambda)}^* \nu^* - \mathcal{Q}_{(\lambda)}^* |\mathbf{q}^*|) \left( 1 + \frac{Q^2}{M^2} - \frac{3W}{M} \right) \frac{G^V(Q^2)}{6 |\mathbf{q}|^2},$$

$$B_{(\lambda)} = B_{(\lambda)}^A = \sqrt{\frac{\Omega}{2}} \left( \mathcal{Q}_{(\lambda)}^* + \nu_{(\lambda)}^* \frac{|\mathbf{q}^*|}{aM} \right) \frac{ZG^A(Q^2)}{3W |\mathbf{q}^*|},$$

$$C_{(\lambda)} = C_{(\lambda)}^A = \left[ (Q_{(\lambda)}^* |\mathbf{q}^*| - \nu_{(\lambda)}^* \nu^*) \left( \frac{1}{3} + \frac{\nu^*}{aM} \right) + \nu_{(\lambda)}^* \left( \frac{2}{3}W - \frac{Q^2}{aM} + \frac{n\Omega}{3aM} \right) \right] \frac{ZG^A(Q^2)}{2W |\mathbf{q}^*|}.$$

Here

$$\nu^* = E_\nu^* - E_\ell^* = \frac{M\nu - Q^2}{W}, \quad a = 1 + \frac{W^2 + Q^2 + M^2}{2MW},$$

$G^{V,A}(Q^2)$  are the vector and axial transition form factors and the remaining notation is explained in Ref. [6]. Other 5 coefficients listed in Eq. (3.11) of Ref. [6] are left unchanged.

The resonance production amplitudes,  $f_\varkappa^\lambda$  with  $\varkappa = \pm 3, \pm 1$  depend on neither final lepton helicity nor initial lepton (neutrino or antineutrino)<sup>5</sup>:

$$[f_\varkappa^\lambda]_\nu = [f_\varkappa^{-\lambda}]_\nu = [f_\varkappa^\lambda]_{\bar{\nu}} = [f_\varkappa^{-\lambda}]_{\bar{\nu}}, \quad \varkappa = \pm 3, \pm 1. \quad (1.53)$$

However, it is not so for  $f_{0\pm}^\lambda$ . Indeed, to find the amplitude  $f_{0\pm}^\lambda$  one need use the values of  $S_{(\lambda)}$ ,  $B_{(\lambda)}$  and  $C_{(\lambda)}$ . Due to (1.38) we have

$$[S_{(\lambda)}]_{\bar{\nu}} = [S_{(-\lambda)}]_\nu, \quad [B_{(\lambda)}]_{\bar{\nu}} = [B_{(-\lambda)}]_\nu, \quad [C_{(\lambda)}]_{\bar{\nu}} = [C_{(-\lambda)}]_\nu. \quad (1.54)$$

Consequently

$$[f_{0\pm}^\lambda]_\nu = [f_{0\pm}^{-\lambda}]_{\bar{\nu}}. \quad (1.55)$$

Using Eqs. (1.53) and (1.55) we obtain

$$[a_\varkappa^\lambda]_\nu = [a_\varkappa^{-\lambda}]_\nu = [a_\varkappa^\lambda]_{\bar{\nu}} = [a_\varkappa^{-\lambda}]_{\bar{\nu}}, \quad \varkappa = \pm 3, \pm 1 \quad (1.56)$$

$$[a_\varkappa^\lambda]_\nu = [a_\varkappa^{-\lambda}]_{\bar{\nu}}, \quad \varkappa = 0\pm, \quad (1.57)$$

and accounting for (1.50)

$$[A_\varkappa^\lambda]_\nu = [A_\varkappa^{-\lambda}]_\nu = [A_\varkappa^\lambda]_{\bar{\nu}} = [A_\varkappa^{-\lambda}]_{\bar{\nu}}, \quad \varkappa = \pm 3, \pm 1$$

$$[A_\varkappa^\lambda]_\nu = [A_\varkappa^{-\lambda}]_{\bar{\nu}}, \quad \varkappa = 0\pm.$$

For the partial cross sections (1.48) the following identities are hold:

$$\begin{aligned} [\sigma_{L,R}^{++}]_\nu &= [\sigma_{L,R}^{+-}]_\nu = [\sigma_{L,R}^{-+}]_\nu = [\sigma_{L,R}^{--}]_\nu = \\ [\sigma_{L,R}^{++}]_{\bar{\nu}} &= [\sigma_{L,R}^{+-}]_{\bar{\nu}} = [\sigma_{L,R}^{-+}]_{\bar{\nu}} = [\sigma_{L,R}^{--}]_{\bar{\nu}}, \\ [\sigma_S^{++}]_\nu &= [\sigma_S^{--}]_{\bar{\nu}}, [\sigma_S^{--}]_\nu = [\sigma_S^{++}]_{\bar{\nu}}, \\ [\sigma_S^{+-}]_\nu &= [\sigma_S^{-+}]_{\bar{\nu}} = [\sigma_S^{+-}]_{\bar{\nu}} = [\sigma_S^{-+}]_{\bar{\nu}}. \end{aligned} \quad (1.58)$$

Next (see Eq. (1.46))

$$\begin{aligned} [\Sigma_{++} + \Sigma_{--}]_\nu &= [c_L^+]^2 [\sigma_L^{++}]_\nu + [c_R^+]^2 [\sigma_R^{++}]_\nu + [c_S^+]^2 [\sigma_S^{++}]_\nu + \\ & \quad [c_L^-]^2 [\sigma_L^{--}]_\nu + [c_R^-]^2 [\sigma_R^{--}]_\nu + [c_S^-]^2 [\sigma_S^{--}]_\nu \end{aligned} \quad (1.59)$$

Using the transition conditions (1.39) and (1.58)

$$[\Sigma_{++} + \Sigma_{--}]_{\bar{\nu}} = [c_L^+]^2 [\sigma_L^{++}]_{\bar{\nu}} + [c_R^+]^2 [\sigma_R^{++}]_{\bar{\nu}} + [c_S^+]^2 [\sigma_S^{++}]_{\bar{\nu}} +$$

<sup>5</sup>The explicit form of the amplitudes can be found in Ref. [6].



$$\begin{aligned}
& [c_L^-]^2 [\sigma_L^{--}]_{\bar{\nu}} + [c_R^-]^2 [\sigma_R^{--}]_{\bar{\nu}} + [c_S^-]^2 [\sigma_S^{--}]_{\bar{\nu}} = \\
& [c_R^-]^2 [\sigma_L^{++}]_{\nu} + [c_L^-]^2 [\sigma_R^{++}]_{\nu} + [c_S^-]^2 [\sigma_S^{--}]_{\nu} + \\
& [c_R^+]^2 [\sigma_L^{--}]_{\nu} + [c_L^+]^2 [\sigma_R^{--}]_{\nu} + [c_S^+]^2 [\sigma_S^{++}]_{\nu}.
\end{aligned} \tag{1.60}$$

Using Mino's "recipe" (1.45) one finds

$$\begin{aligned}
[\Sigma_{++} + \Sigma_{--}]_{\bar{\nu}} &= [c_L^+]^2 [\sigma_L^{++}]_{\nu} + [c_R^+]^2 [\sigma_R^{++}]_{\nu} + [c_S^+]^2 [\sigma_S^{++}]_{\nu} + \\
& [c_L^-]^2 [\sigma_L^{--}]_{\nu} + [c_R^-]^2 [\sigma_R^{--}]_{\nu} + [c_S^-]^2 [\sigma_S^{--}]_{\nu} = \\
& [c_R^+]^2 [\sigma_L^{++}]_{\nu} + [c_L^+]^2 [\sigma_R^{++}]_{\nu} + [c_S^+]^2 [\sigma_S^{++}]_{\nu} + \\
& [c_R^-]^2 [\sigma_L^{--}]_{\nu} + [c_L^-]^2 [\sigma_R^{--}]_{\nu} + [c_S^-]^2 [\sigma_S^{--}]_{\nu},
\end{aligned} \tag{1.61}$$

what at first glance is different from Eq. (1.60), but again using the relations (1.58) we obtain

$$\begin{aligned}
[\Sigma_{++} + \Sigma_{--}]_{\bar{\nu}} &= [c_R^+]^2 [\sigma_L^{++}]_{\nu} + [c_L^+]^2 [\sigma_R^{++}]_{\nu} + [c_S^+]^2 [\sigma_S^{++}]_{\nu} + \\
& [c_R^-]^2 [\sigma_L^{--}]_{\nu} + [c_L^-]^2 [\sigma_R^{--}]_{\nu} + [c_S^-]^2 [\sigma_S^{--}]_{\nu} = \\
& [c_R^+]^2 [\sigma_L^{--}]_{\nu} + [c_L^+]^2 [\sigma_R^{--}]_{\nu} + [c_S^+]^2 [\sigma_S^{--}]_{\nu} + \\
& [c_R^-]^2 [\sigma_L^{++}]_{\nu} + [c_L^-]^2 [\sigma_R^{++}]_{\nu} + [c_S^-]^2 [\sigma_S^{++}]_{\nu}.
\end{aligned} \tag{1.62}$$

It can be seen that Eqs. (1.60) and (1.62) are coincide. Thus differential cross section of unpolarized lepton production (1.47) does not depend on what transition conditions is used.

Now, by applying transition conditions (1.39) and (1.58) we get:

$$\begin{aligned}
[\Sigma_{\lambda\lambda'}]_{\bar{\nu}} &= [c_L^\lambda]_{\bar{\nu}} [c_L^{\lambda'}]_{\bar{\nu}} [\sigma_L^{\lambda\lambda'}]_{\bar{\nu}} + [c_R^\lambda]_{\bar{\nu}} [c_R^{\lambda'}]_{\bar{\nu}} [\sigma_R^{\lambda\lambda'}]_{\bar{\nu}} + [c_S^\lambda]_{\bar{\nu}} [c_S^{\lambda'}]_{\bar{\nu}} [\sigma_S^{\lambda\lambda'}]_{\bar{\nu}} = \\
& \lambda\lambda' \left( [c_R^{-\lambda}]_{\nu} [c_R^{-\lambda'}]_{\nu} [\sigma_L^{\lambda\lambda'}]_{\nu} + [c_L^{-\lambda}]_{\nu} [c_L^{-\lambda'}]_{\nu} [\sigma_R^{\lambda\lambda'}]_{\nu} + [c_S^{-\lambda}]_{\nu} [c_S^{-\lambda'}]_{\nu} [\sigma_S^{\lambda-\lambda'}]_{\nu} \right).
\end{aligned}$$

But, according to Eqs. (1.45), Mino's "recipe" gives different (in fact wrong) result:

$$\begin{aligned}
[\Sigma_{\lambda\lambda'}]_{\bar{\nu}} &= [c_L^\lambda]_{\bar{\nu}} [c_L^{\lambda'}]_{\bar{\nu}} [\sigma_L^{\lambda\lambda'}]_{\nu} + [c_R^\lambda]_{\bar{\nu}} [c_R^{\lambda'}]_{\bar{\nu}} [\sigma_R^{\lambda\lambda'}]_{\nu} + [c_S^\lambda]_{\bar{\nu}} [c_S^{\lambda'}]_{\bar{\nu}} [\sigma_S^{\lambda\lambda'}]_{\nu} = \\
& [c_R^\lambda]_{\nu} [c_R^{\lambda'}]_{\nu} [\sigma_L^{\lambda\lambda'}]_{\nu} + [c_L^\lambda]_{\nu} [c_L^{\lambda'}]_{\nu} [\sigma_R^{\lambda\lambda'}]_{\nu} + [c_S^\lambda]_{\nu} [c_S^{\lambda'}]_{\nu} [\sigma_S^{\lambda\lambda'}]_{\nu}.
\end{aligned} \tag{1.63}$$

Now let us consider the case of the  $\nu \mapsto \bar{\nu}$  transition for the differential cross section of single pion production derived in Refs. [2, 3]:

$$\begin{aligned}
\frac{d\sigma(\nu N \rightarrow lN\pi)}{dQ^2 dW d\Omega_\pi} &= \frac{G_F^2}{2} \frac{1}{(2\pi)^4} \frac{|\mathbf{q}|}{4} \frac{Q^2}{(k^L)^2} \sum_{\lambda_2, \lambda_1} \left\{ \right. \\
& \left| C_{L-} (\tilde{F}_{\lambda_2 \lambda_1}^L(\theta, \phi) - \tilde{G}_{\lambda_2 \lambda_1}^L(\theta, \phi)) + C_{R-} (\tilde{F}_{\lambda_2 \lambda_1}^R(\theta, \phi) - \tilde{G}_{\lambda_2 \lambda_1}^R(\theta, \phi)) + \right. \\
& \left. C_- (\tilde{F}_{\lambda_2 \lambda_1}^-(\theta, \phi) - \tilde{G}_{\lambda_2 \lambda_1}^-(\theta, \phi)) \right|^2 \\
& + \left| C_{L+} (\tilde{F}_{\lambda_2 \lambda_1}^L(\theta, \phi) - \tilde{G}_{\lambda_2 \lambda_1}^L(\theta, \phi)) + C_{R+} (\tilde{F}_{\lambda_2 \lambda_1}^R(\theta, \phi) - \tilde{G}_{\lambda_2 \lambda_1}^R(\theta, \phi)) + \right. \\
& \left. C_+ (\tilde{F}_{\lambda_2 \lambda_1}^+(\theta, \phi) - \tilde{G}_{\lambda_2 \lambda_1}^+(\theta, \phi)) \right|^2 \left. \right\}.
\end{aligned} \tag{1.64}$$

First, we suppose, that the amplitudes  $\tilde{F}_{\lambda_2\lambda_1}^{\lambda_k}(\theta, \phi)$  and  $\tilde{G}_{\lambda_2\lambda_1}^{\lambda_k}(\theta, \phi)$  correspond only to the resonance case. Using formulas from Table 1.1 and transition relations (1.53) and (1.55), we can find how the amplitudes should be transformed when we want to calculate cross sections for antineutrino:

$$\begin{aligned} \left[ \tilde{F}_{\lambda_2\lambda_1}^L \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^L \right]_{\bar{\nu}} &= \left[ \tilde{F}_{\lambda_2\lambda_1}^L \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^L \right]_{\nu}, & \left[ \tilde{F}_{\lambda_2\lambda_1}^R \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^R \right]_{\bar{\nu}} &= \left[ \tilde{F}_{\lambda_2\lambda_1}^R \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^R \right]_{\nu}, \\ \left[ \tilde{F}_{\lambda_2\lambda_1}^+ \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^+ \right]_{\bar{\nu}} &= \left[ \tilde{F}_{\lambda_2\lambda_1}^- \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^- \right]_{\nu}, & \left[ \tilde{F}_{\lambda_2\lambda_1}^- \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^- \right]_{\bar{\nu}} &= \left[ \tilde{F}_{\lambda_2\lambda_1}^+ \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^+ \right]_{\nu}. \end{aligned} \quad (1.65)$$

For example, it is easy to check that:

$$\begin{aligned} \left[ \tilde{F}_{\frac{1}{2}\frac{1}{2}}^L \right]_{\bar{\nu}} - \left[ \tilde{G}_{\frac{1}{2}\frac{1}{2}}^L \right]_{\bar{\nu}} &= \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) [f_{+3}^\lambda]_{\bar{\nu}}(R) d_{\frac{3}{2}\frac{1}{2}}^j(\theta) e^{-2i\phi} = \\ &= \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) [f_{+3}^\lambda]_{\nu}(R) d_{\frac{3}{2}\frac{1}{2}}^j(\theta) e^{-2i\phi} = \left[ \tilde{F}_{\frac{1}{2}\frac{1}{2}}^L \right]_{\nu} - \left[ \tilde{G}_{\frac{1}{2}\frac{1}{2}}^L \right]_{\nu}, \\ \left[ \tilde{F}_{\frac{1}{2}\frac{1}{2}}^- \right]_{\bar{\nu}} - \left[ \tilde{G}_{\frac{1}{2}\frac{1}{2}}^- \right]_{\bar{\nu}} &= + \frac{|\mathbf{k}|}{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) [f_{0-}^{(-)}]_{\bar{\nu}}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{-i\phi} = \\ &= + \frac{|\mathbf{k}|}{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) [f_{0-}^{(+)}]_{\nu}(R) d_{\frac{1}{2}\frac{1}{2}}^j(\theta) e^{-i\phi} = \left[ \tilde{F}_{\frac{1}{2}\frac{1}{2}}^+ \right]_{\nu} - \left[ \tilde{G}_{\frac{1}{2}\frac{1}{2}}^+ \right]_{\nu}, \end{aligned}$$

etc. Applying the transition relations (1.43) to Eq. (1.64) one gets:

$$\begin{aligned} \frac{d\sigma(\bar{\nu}N \rightarrow lN\pi)}{dQ^2 dW d\Omega_\pi} &= \frac{G_F^2}{2} \frac{1}{(2\pi)^4} \frac{|\mathbf{q}|}{4} \frac{Q^2}{(k^L)^2} \sum_{\lambda_2, \lambda_1} \left\{ \right. \\ & \left| [C_{L-}]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\bar{\nu}} \right) + [C_{R-}]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\bar{\nu}} \right) \right. \\ & \left. [C_{-}]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^-(\theta, \phi) \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^-(\theta, \phi) \right]_{\bar{\nu}} \right) \right|^2 + \\ & \left| [C_{L+}]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\bar{\nu}} \right) + [C_{R+}]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\bar{\nu}} \right) \right. \\ & \left. [C_{+}]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^+(\theta, \phi) \right]_{\bar{\nu}} - \left[ \tilde{G}_{\lambda_2\lambda_1}^+(\theta, \phi) \right]_{\bar{\nu}} \right) \right|^2 \left. \right\} = \\ & \frac{G_F^2}{2} \frac{1}{(2\pi)^4} \frac{|\mathbf{q}|}{4} \frac{Q^2}{(k^L)^2} \sum_{\lambda_2, \lambda_1} \left\{ \right. \\ & \left| - [C_{R+}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\nu} \right) - [C_{L+}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\nu} \right) \right. \\ & \left. [C_{+}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^+(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^+(\theta, \phi) \right]_{\nu} \right) \right|^2 + \\ & \left| [C_{R-}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\nu} \right) + [C_{L-}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\nu} \right) \right. \\ & \left. [C_{-}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^-(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^-(\theta, \phi) \right]_{\nu} \right) \right|^2 \left. \right\} = \\ & \frac{G_F^2}{2} \frac{1}{(2\pi)^4} \frac{|\mathbf{q}|}{4} \frac{Q^2}{(k^L)^2} \sum_{\lambda_2, \lambda_1} \left\{ \right. \\ & \left| [C_{R+}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^L(\theta, \phi) \right]_{\nu} \right) + [C_{L+}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^R(\theta, \phi) \right]_{\nu} \right) \right. \\ & \left. [C_{+}]_{\nu} \left( \left[ \tilde{F}_{\lambda_2\lambda_1}^+(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2\lambda_1}^+(\theta, \phi) \right]_{\nu} \right) \right|^2 \left. \right\} = \end{aligned}$$

$$\begin{aligned}
& \left[ C_{+} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^{+}(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^{+}(\theta, \phi) \right]_{\nu} \right)^2 + \\
& \left[ C_{R-} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} \right) + \left[ C_{L-} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} \right) - \\
& \left[ C_{-} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^{-}(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^{-}(\theta, \phi) \right]_{\nu} \right)^2 \left. \right\}. \tag{1.66}
\end{aligned}$$

If we use the transition relations (1.44), we get

$$\begin{aligned}
\frac{d\sigma(\bar{\nu}N \rightarrow lN\pi)}{dQ^2 dW d\Omega_{\pi}} &= \frac{G_F^2}{2} \frac{1}{(2\pi)^4} \frac{|\mathbf{q}|}{4} \frac{Q^2}{(k^L)^2} \sum_{\lambda_2, \lambda_1} \left\{ \right. \\
& \left[ C_{L-} \right]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} \right) + \left[ C_{R-} \right]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} \right) + \\
& \left[ C_{-} \right]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^{-}(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^{-}(\theta, \phi) \right]_{\nu} \right)^2 + \\
& \left[ C_{L+} \right]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} \right) + \left[ C_{R+} \right]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} \right) + \\
& \left. \left[ C_{+} \right]_{\bar{\nu}} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^{+}(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^{+}(\theta, \phi) \right]_{\nu} \right)^2 \right\} = \\
& \frac{G_F^2}{2} \frac{1}{(2\pi)^4} \frac{|\mathbf{q}|}{4} \frac{Q^2}{(k^L)^2} \sum_{\lambda_2, \lambda_1} \left\{ \right. \\
& \left[ C_{R-} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} \right) + \left[ C_{L-} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} \right) + \\
& \left[ C_{-} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^{-}(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^{-}(\theta, \phi) \right]_{\nu} \right)^2 + \\
& \left[ C_{R+} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^L(\theta, \phi) \right]_{\nu} \right) + \left[ C_{L+} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^R(\theta, \phi) \right]_{\nu} \right) + \\
& \left. \left[ C_{+} \right]_{\nu} \left( \left[ \tilde{F}_{\lambda_2 \lambda_1}^{+}(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^{+}(\theta, \phi) \right]_{\nu} \right)^2 \right\}. \tag{1.67}
\end{aligned}$$

We see that Eqs. (1.66) and (1.67) differ in sign of the term  $\left[ \tilde{F}_{\lambda_2 \lambda_1}^{\pm}(\theta, \phi) \right]_{\nu} - \left[ \tilde{G}_{\lambda_2 \lambda_1}^{\pm}(\theta, \phi) \right]_{\nu}$ .

## Conclusions:

Ultimately, Minoo's "recipe" for the  $\nu \mapsto \bar{\nu}$  transition gives correct result only for the double differential cross sections, like  $d\sigma/dWdQ^2$ . For the cross section  $d\sigma/dQ^2dWd\Omega_{\pi}$  and for the polarization density matrix her "recipe" does not work.

Let's emphasize that calculation of the polarization matrix is by no means an academic issue because in the long-awaited high-precision experiments with atmospheric and astrophysical neutrinos (like Hyper-Kamiokande, DUNE, PINGU, ORCA) and in possible future dedicated accelerator experiments, one will need to perform detailed calculations of production and decay of polarized  $\tau$ -leptons in the detector, Earth, atmosphere, and astrophysical sources. In fact, this was one of the main goals in generalizing the RS model (KLN, BS, etc.).<sup>6</sup> We think, this option will certainly need to be added to GENIE. So we plan to return to this problem in the future.

Considering that the transition rules (1.39) are consequence of the leptonic current properties only, they are valid for any neutrino-induced CC process. The coincidence of the cross

<sup>6</sup>More details can be found in Refs. [8, 10].

section  $d\sigma/dWdQ^2$  obtained by using this rules with Minoo's one is due to underlying symmetry of the Rein-Sehgal model [6]. This may not be so in other models or after some modifications of the KLN-BS or MK models. In particular, since in the MK-model, the resonance amplitudes are coherently added to the non-resonance ones, it is not in general obvious that in this case even the cross section  $d\sigma/dWdQ^2$  calculated by using Minoo's prescription will remain correct.

Let us also note that agreement of the MK model predictions with the experimental data cannot confirm or disconfirm the  $\nu \mapsto \bar{\nu}$  transition rules just because the currently available data are too fragmentary and uncertain and are either slowly sensible or (as the single and double-differential cross sections) fully insensible to the differences in the two versions of the rules under consideration. Moreover, after refitting the MK model parameters, the agreement with the more detailed data may seem to be quite satisfactory even with wrong formulas. But the predictive power of the model will be under big question. So we suggest to use the  $\nu \mapsto \bar{\nu}$  transition rules derived in Ref. [8] and reproduced above... or to refute our derivations.

# Chapter 2

## Fixed problems, bugs and mistakes

### 2.1 Incorrect phase factors for resonance and background amplitudes

The expression for the differential cross section is obtained assuming incorrect phase factors for resonance and background amplitudes in the C++ code provided by Minoo. The proof of this fact can be founded in accompanying *Wolfram Mathematica* notebook **XSec.MK.Diff.nb** with all explanatory comments. In brief, the phase factors of the amplitudes  $\tilde{F}_{\lambda_2, \lambda_1}^{eL}(\theta, \phi)$  and  $\tilde{F}_{\lambda_2, \lambda_1}^{eR}(\theta, \phi)$  are mixed up. This lead to incorrect expression for the cross section.

There is a similar problem in the latest version of Minoo's code. We checked differential cross section only for one channel (*MK\_mode1.cc*) and found that it is wrong (see accompanying file **XSec.MK.Diff.v.2.nb**), but we didn't check other channels yet.

### 2.2 Incorrect helicity amplitudes

The helicity amplitudes:

- for resonance  $S_{11}(1650)$  instead of

$$f1 = \text{sqrt}(1./24.)*L*(R1\_minus + 4*\text{sin}2\text{Wein}*R1\_V);$$

should be

$$f1 = \text{sqrt}(1./24.)*L*(R1\_plus + 4*\text{sin}2\text{Wein}*R1\_V);$$

- for resonance  $P_{13}(1720)$  instead of

$$\begin{aligned} f\_1 &= -\text{sqrt}(27./40.)*L*T2\_minus + \\ &\quad \text{sqrt}(5./12.)*L*L*(R2\_minus+2.*\text{sin}2\text{Wein}*(4./5.)*R2\_V); \\ f1 &= +\text{sqrt}(27./40.)*L*T2\_plus - \\ &\quad \text{sqrt}(5./12.)*L*L*(R2\_plus + 2.*\text{sin}2\text{Wein}*(4./5.)*R2\_V); \end{aligned}$$

```

f3      = -sqrt(9./40.)*L*T2_plus;
f0_plus = -sqrt(3./20.)*L*L*S2_KLM_minus +
          sqrt(5./12.)*L*(L*C2_minus-5.*B2_minus);
f0_minus = sqrt(3./20.)*L*L*S2_KLM_minus +
          sqrt(5./12.)*L*(L*C2_minus-5.*B2_minus);

```

should be

```

f_1      = sqrt(27./40.)*L*T2_minus +
          sqrt(5./12.)*L*L*(R2_minus+2.*sin2Wein*(4./5.)*R2_V);
f1       = -sqrt(27./40.)*L*T2_plus -
          sqrt(5./12.)*L*L*(R2_plus +2.*sin2Wein*(4./5.)*R2_V);
f3       = sqrt(9./40.)*L*T2_plus;
f0_plus  = sqrt(3./20.)*L*L*S2_KLM_minus +
          sqrt(5./12.)*L*(L*C2_minus-5.*B2_minus);
f0_minus = -sqrt(3./20.)*L*L*S2_KLM_minus +
          sqrt(5./12.)*L*(L*C2_minus-5.*B2_minus);

```

- for resonance  $F_{15}(1680)$  instead of

```

f3      = sqrt(9./10.)*L*T2_plus;

```

should be

```

f3      = sqrt(9./10.)*L*T2_plus;

```

defined in the file *MK\_imode7.cc* (latest code version) are wrong (see Ref. [5]).

## 2.3 Erroneous dynamical form factor B

In the MK-model the wrong form factor  $B$  was used (see Minoos erratum to Ref. [10]). It was fixed in the file *mk\_imode1\_new.cc* and *mk\_imode4\_new.cc*.

## 2.4 Mistaken cross-section formula for NC-processes in the code

The values `Fem_zero_minus` is not used in the expression for the cross section. As a result some parts of the expression for the cross section are incorrect, for example, instead of lines 1003-1006 from *mk\_imode4\_new.cc*:

```

+ pow( ((1. - 2.*sin2Wein)*F_zero_minus11 - G_zero_minus11 + C3*sum3Re_OM11 + C1*sum1Re_OM11) , 2)
+ pow( ((1. - 2.*sin2Wein)*F_zero_minus_11 - G_zero_minus_11 + C3*sum3Re_OM_11 + C1*sum1Re_OM_11) , 2)
+ pow( ((1. - 2.*sin2Wein)*F_zero_minus1_1 - G_zero_minus1_1 + C3*sum3Re_OM1_1 + C1*sum1Re_OM1_1) , 2)
+ pow( ((1. - 2.*sin2Wein)*F_zero_minus_1_1 - G_zero_minus_1_1 + C3*sum3Re_OM_1_1+ C1*sum1Re_OM_1_1) , 2)

```

should be the following lines:

```

+ pow( ((1. - 2.*sin2Wein)*F_zero_minus11 - G_zero_minus11 - 4.*sin2Wein*Fem_zero_minus11
+ C3*sum3Re_OM11 + C1*sum1Re_OM11 ) , 2)
+ pow( ((1. - 2.*sin2Wein)*F_zero_minus_11 - G_zero_minus_11 - 4.*sin2Wein*Fem_zero_minus_11
+ C3*sum3Re_OM_11 + C1*sum1Re_OM_11) , 2)
+ pow( ((1. - 2.*sin2Wein)*F_zero_minus1_1 - G_zero_minus1_1 - 4.*sin2Wein*Fem_zero_minus1_1
+ C3*sum3Re_OM1_1 + C1*sum1Re_OM1_1) , 2)
+ pow( ((1. - 2.*sin2Wein)*F_zero_minus_1_1 - G_zero_minus_1_1- 4.*sin2Wein*Fem_zero_minus_1_1
+ C3*sum3Re_OM_1_1+ C1*sum1Re_OM_1_1), 2)

```

The error has been fixed in the most recent version of the file: *mk\_imode4\_new.cc*.

## 2.5 Incorrect $C_j$ -signs

The  $C_j$ -signs in the following fragments of code, which are denoted as Jsgn, are wrong, because for the amplitudes  $\tilde{F}_{-\frac{1}{2},\lambda_1}^{eR}(\theta, \phi)$  (denoted as HV\_(real|Im)P\_1(1|\_1)) the corresponding signs are always positive (see Table 7 from Ref. [5]).

```

const int Jsgn[nRes] = {1,-1,-1, 1, 1, 1, 1,-1,-1,-1,-1, 1,-1,-1, 1, 1, 1};
HV_realP11[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_1;
HV_realP_11[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_1*Jsgn[i];
HV_realP1_1[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_3;
HV_realP_1_1[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_3*Jsgn[i];

HV_ImP11[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_1;
HV_ImP_11[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_1*Jsgn[i];
HV_ImP1_1[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_3;
HV_ImP_1_1[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_3*Jsgn[i];

HA_realP11[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_1;
HA_realP_11[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_1*Jsgn[i];
HA_realP1_1[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_3;
HA_realP_1_1[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_real[i]*fV_3*Jsgn[i];

HA_ImP11[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_1;
HA_ImP_11[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_1*Jsgn[i];
HA_ImP1_1[i] = -sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_3;
HA_ImP_1_1[i] = sqrt(2.)*JP[i]*Dsgn[i]*kapa[i]*Vf_BW_Im[i]*fV_3*Jsgn[i];

```

It was fixed in the file *mk\_imode1\_erratum.cc*.

## 2.6 Problem with multipole expansion of some amplitudes in the code

The multipole expansion of the following amplitudes  $\tilde{F}_{\text{res}-\frac{1}{2},\frac{1}{2}}^+$  and  $\tilde{F}_{\text{res}-\frac{1}{2},\frac{1}{2}}^-$  is definitely incorrect as they were defined in the first version of Erratum (this is not the same definition as in paper [3]):

$$\begin{aligned}\tilde{F}_{\text{res}-\frac{1}{2},\frac{1}{2}}^- &= \frac{|\mathbf{k}|}{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0-}^{V(-)}(R) d_{-\frac{1}{2},\frac{1}{2}}^j(\theta), \\ \tilde{F}_{\text{res}-\frac{1}{2},\frac{1}{2}}^+ &= \frac{|\mathbf{k}|}{\sqrt{Q^2}} \sum_j \frac{2j+1}{\sqrt{2}} \mathcal{D}^j(R) f_{0-}^{V(-)}(R) d_{-\frac{1}{2},\frac{1}{2}}^j(\theta).\end{aligned}\tag{2.1}$$

But in the MK-code they are defined different way. For example, in line 823 of the file *mk\_imode1\_new.cc*:

```
double sum3Re_OM_11 =  HA0_realM_11[0] *d31_1 + HA0_realM_11[4]*d11_1 +
                        HA0_realM_11[9] *d31_1 + HA0_realM_11[12]*d51_1+
                        HA0_realM_11[13]*d11_1 + HA0_realM_11[14]*d31_1+
                        HA0_realM_11[15]*d71_1 + HA0_realM_11[16]*d31_1;
```

where  $d_{j1_1}$  denotes  $d_{\frac{1}{2},-\frac{1}{2}}^j$ . But, according to Eq. (1.16),  $d_{\frac{1}{2},-\frac{1}{2}}^j = -d_{-\frac{1}{2},\frac{1}{2}}^j$ . So the above line should be rewritten as follows:

```
double sum3Re_OM_11 = -(HA0_realM_11[0] *d31_1 + HA0_realM_11[4]*d11_1 +
                        HA0_realM_11[9] *d31_1 + HA0_realM_11[12]*d51_1+
                        HA0_realM_11[13]*d11_1 + HA0_realM_11[14]*d31_1+
                        HA0_realM_11[15]*d71_1 + HA0_realM_11[16]*d31_1);
```

Lines 828, 843, 848, 864, 869, 884, and 889 should be corrected in the same way.

The mistake has been fixed in the second version of Erratum [4] but appears again in the latest versions of erratum for paper [3] and code. However, it is unclear how one should fix it (see section 1.3.4).

## 2.7 Expressions for EM-amplitudes contains mistake

There is an error in the lines 207–210 of the file *mk\_imode4\_new.cc*, instead of *C\_S\_plus\_square* it should be *C\_S\_minus\_square*:

```
double Fem_zero_minus11 = (1./sqrt(C_S_plus_square))*sqrt((1+t)/2.)*(k_0*eps_z_L - abs_mom_k*eps_zero_L)*(sFem_5 + sFem_6);
double Fem_zero_minus_11 = -(1./sqrt(C_S_plus_square))*sqrt((1-t)/2.)*(k_0*eps_z_L - abs_mom_k*eps_zero_L)*(sFem_5 - sFem_6);
double Fem_zero_minus1_1 = -(1./sqrt(C_S_plus_square))*sqrt((1-t)/2.)*(k_0*eps_z_L - abs_mom_k*eps_zero_L)*(sFem_5 - sFem_6);
double Fem_zero_minus_1_1 = -(1./sqrt(C_S_plus_square))*sqrt((1+t)/2.)*(k_0*eps_z_L - abs_mom_k*eps_zero_L)*(sFem_5 + sFem_6);
```

## 2.8 Error in the dynamical amplitudes for resonance $D_{13}(1520)$

The error is in lines 448, 449 of the file *mk\_imode6.cc*:



```
fV_1 = -sqrt(3./2.)*(-T1_V[2]+2.*sin2Wein*T1_V[2])+sqrt(4./3.)*L*(-R1_V[2]+3.*sin2Wein*R1_V[2]);
fV1 = -sqrt(3./2.)*(-T1_V[2]+2.*sin2Wein*T1_V[2])+sqrt(4./3.)*L*(-R1_V[2]+3.*sin2Wein*R1_V[2]);
```

The corrected lines should be [6]:

```
fV_1 = -sqrt(3./2.)*(-T1_V[2]+2.*sin2Wein*T1_V[2])+sqrt(4./3.)*L*(-R1_V[2]+sin2Wein*R1_V[2]);
fV1 = -sqrt(3./2.)*(-T1_V[2]+2.*sin2Wein*T1_V[2])+sqrt(4./3.)*L*(-R1_V[2]+sin2Wein*R1_V[2]);
```

## 2.9 Wrong sign for As-values in the code

The error is in lines 263–268 of *mk\_imode6.cc* and *mk\_imode16.cc*:

```
double As_1=
  -(1./2.1./sqrt(2.))*(g_A/f_pi)*Gs_A*FA_cut*(2.*M/(m_pi*m_pi - 2.*q_0*p_10
  - 2.*abs_mom_q*abs_mom_k*t))-sqrt(2.)*(g_A/f_pi)*Gs_A*FA_cut*(M/(W*W - M*M)));
```

The corrected lines in all files *mk\_imode4.cc*, *mk\_imode5.cc*, *mk\_imode7.cc*, *mk\_imode14.cc*, *mk\_imode15.cc* and *mk\_imode17.cc* for other channels are:

```
double As_1=
  (1./2.1./sqrt(2.))*(g_A/f_pi)*Gs_A*FA_cut*(2.*M/(m_pi*m_pi - 2.*q_0*p_10
  - 2.*abs_mom_q*abs_mom_k*t))+sqrt(2.)*(g_A/f_pi)*Gs_A*FA_cut*(M/(W*W - M*M)));
```

which follows from isospin symmetry.

The problem remains in the latest version of the code.

## 2.10 Mistype in a formula for cross section

A mistype is in line 1201 of the file *mk\_imode2.cc*, namely, the amplitude `sum1Re0PMM` should be instead of `sum1Re0MMM`. For more details see the accompanying *Wolfram Mathematica* notebook `XSec_MK_Diff.nb`.

## 2.11 Problem with phase factors for resonance amplitudes

The phase factors  $\exp[n\phi]$  of the resonance amplitudes in Table 3 in Ref. [3] are incorrect. They should be the same as for the background contribution presented in Table 6 of the same paper. This mistake has been fixed in Ref. [4].

# Bibliography

- [1] M. Kabirnezhad. *Improvement of Single Pion Production for T2K experiment simulation tools*. PhD thesis, National Center for Nuclear Research, Otwock-Świerk, 2017. URL: <s3.cern.ch/inspire-prod-files-a/a29746a5bc5392dda0220c658228916f>.
- [2] M. Kabirnezhad. *Improvement of Single Pion Production for T2K experiment simulation tools(update)*. PhD thesis, National Center for Nuclear Research, Otwock-Świerk, 2020. URL: <ncbj.gov.pl/sites/default/files/m.kabirnezhad-thesis\_final\_0.pdf>.
- [3] M. Kabirnezhad. Single pion production in neutrino-nucleon Interactions. *Phys. Rev. D*, 97:013002, 2018. [Erratum: arXiv:1711.02403].
- [4] M. Kabirnezhad. Single pion production in neutrino-nucleon Interactions. *Erratum (private communication, December 4, 2019)*.
- [5] D. Rein. Angular distribution in neutrino-induced single pion production processes. *Zeitschrift für Physik C Particles and Fields*, 35:43–64, 1987.
- [6] D. Rein and L.M. Sehgal. Neutrino Excitation of Baryon Resonances and Single Pion Production. *Annals Phys.*, 133:79–153, 1981.
- [7] M. Jacob and G.C. Wick. On the general theory of collisions for particles with spin. *Annals of Physics*, 7:404–428, 1959.
- [8] K.S. Kuzmin, V.V. Lyubushkin, and V.A. Naumov. Lepton polarization in neutrino nucleon interactions. *Mod. Phys. Lett. A*, 19:2815–2829, 2004.
- [9] F. Ravndal. Weak production of nuclear resonances in a relativistic quark model. *Nuovo Cim. A*, 18:385–415, 1973.
- [10] Ch. Berger and L.M. Sehgal. Lepton mass effects in single pion production by neutrinos. *Phys. Rev. D*, 76:113004, 2007. [Erratum: *ibid.*, D, 77, 059901(E) (2008)].